

Discrete Optimization for Image Analysis

Bjoern Andres

University of Tübingen

Outline

- ▶ Quadratic multi-linear polynomial forms and quadratic posiforms
- ▶ Complementation
- ▶ Excursus: Maximum st -Flow and Minimum st -Cut
 - ▶ Definitions
 - ▶ Maximum st -Flow Problem
 - ▶ Residual networks and augmenting paths
 - ▶ Minimum st -Cut Problem
 - ▶ Maximum st -Flow/Minimum st -Cut Theorem
 - ▶ Ford-Fulkerson-Algorithm
- ▶ Strong persistency

For any $n \in \mathbb{N}$, consider n -variate **quadratic** forms:

- ▶ any **multi-linear polynomial form** $c \in C_{n2}$ and $f_c : \{0, 1\}^2 \rightarrow \mathbb{R}$, i.e., for all $x \in \{0, 1\}^n$,

$$f_c(x) = c_\emptyset + \sum_{j \in [n]} c_{\{j\}} x_j + \sum_{\{j,k\} \in \binom{[n]}{2}} c_{\{j,k\}} x_j x_k$$

- ▶ any **posiform** $c' \in C_{n2}^+$ and $f'_c : \{0, 1\}^2 \rightarrow \mathbb{R}$, i.e., for all $x \in \{0, 1\}^n$,

$$\begin{aligned} f'_{c'}(x) &= c'_{\emptyset\emptyset} + \sum_{j \in [n]} \left(c'_{\{j\}\emptyset} x_j + c'_{\emptyset\{j\}} (1 - x_j) \right) \\ &+ \sum_{\{j,k\} \in \binom{[n]}{2}} \left(c'_{\{j,k\}\emptyset} x_j x_k + c'_{\{j\}\{k\}} x_j (1 - x_k) \right. \\ &\quad \left. + c'_{\{k\}\{j\}} x_k (1 - x_j) + c'_{\emptyset\{j,k\}} (1 - x_j)(1 - x_k) \right) \end{aligned}$$

Lemma 1

For any $n \in \mathbb{N}$, any QPBF $f : \{0, 1\}^n \rightarrow \mathbb{R}$, the $c \in C_{n2}$ such that $f_c = f$ and any $c' \in C_{n2}^+(f)$ holds

$$c_{\emptyset} = c'_{\emptyset\emptyset} + \sum_{j=1}^n c'_{\emptyset\{j\}} + \sum_{\{j,k\} \in \binom{[n]}{2}} c'_{\emptyset\{j,k\}}$$

$$\forall j \in [n] \quad c_{\{j\}} = c'_{\{j\}\emptyset} - c'_{\emptyset\{j\}} + \sum_{k \in [n] - \{j\}} (c'_{\{j\}\{k\}} - c'_{\emptyset\{j,k\}})$$

$$\forall \{j,k\} \in \binom{[n]}{2} \quad c_{\{j,k\}} = c'_{\{j,k\}\emptyset} + c'_{\emptyset\{j,k\}} - c'_{\{j\}\{k\}} - c'_{\{k\}\{j\}}$$

Proof.

- ▶ Expansion of the posiform c' yields a quadratic multi-linear polynomial form.
- ▶ Comparison with c yields the conditions stated in the Lemma.

Definition 1 (Complementation)

For any $n \in \mathbb{N}$ and any QPBF $f : \{0, 1\}^n \rightarrow \mathbb{R}$,

$$r_f := \max_{c' \in C_{n^2}^+(f)} c'_{\emptyset\emptyset} \quad (1)$$

is called the **floor dual** of f .

Lemma 2

For any $n \in \mathbb{N}$ and any QPBF $f : \{0, 1\}^n \rightarrow \mathbb{R}$, the floor dual can be computed in polynomial time.

Proof. For the multi-linear polynomial form $c \in C_{n2}$ such that $f_c = f$, r_f is the solution of the linear programming problem below (by Lemma 1).

$$\max_{c': J_{n2}^+ \rightarrow \mathbb{R}} \quad c_\emptyset - \sum_{j=1}^n c'_{\emptyset\{j\}} - \sum_{\{j,k\} \in \binom{[n]}{2}} c'_{\emptyset\{j,k\}}$$

$$\text{subject to} \quad \forall j \in [n] \quad c_{\{j\}} = c'_{\{j\}\emptyset} - c'_{\emptyset\{j\}} + \sum_{k \in [n] - \{j\}} (c'_{\{j\}\{k\}} - c'_{\emptyset\{j,k\}})$$

$$\forall \{j, k\} \in \binom{[n]}{2} \quad c_{\{j,k\}} = c'_{\{j,k\}\emptyset} + c'_{\emptyset\{j,k\}} - c'_{\{j\}\{k\}} - c'_{\{k\}\{j\}}$$

$$\forall J \in J_{n2}^+ - \{(\emptyset, \emptyset)\} \quad 0 \leq c'_J \quad .$$

Can the floor dual be computed more efficiently than by an algorithm for general LPs?

Excursus: Maximum st -Flow and Minimum st -Cut



- ▶ Definitions
- ▶ Maximum st -Flow Problem
- ▶ Residual networks and augmenting paths
- ▶ Minimum st -Cut Problem
- ▶ Maximum st -Flow/Minimum st -Cut Theorem
- ▶ Ford-Fulkerson-Algorithm

Definition 2

A pair (V, E) is called

- ▶ an **undirected graph** iff $E \subseteq \binom{V}{2}$
- ▶ a **directed graph** iff $E \subseteq V^2$.

The elements of V are called **nodes**. The elements of E are called **edges**.



Definition 3


In any directed graph (V, E) ,

- ▶ an edge $e \in E$ is called a **self-edge** iff

$$\exists v \in V \quad e = vv .$$

- ▶ a pair $ee' \in E^2$ of edges is called a **digon** iff

$$\exists v, v' \in V \quad v \neq v' \wedge e = vv' \wedge e' = v'v .$$




Below, the term **directed graph** shall always mean directed graph **without self-edges** and **without digons**, i.e., a pair (V, E) such that $E \subseteq V^2$ and

$$\forall vw \in V^2 \quad vw \notin E \vee wv \notin E .$$

For any directed graph (V, E) , any $U \subseteq V$ and any $W \subseteq V$ let

$$UW := \{uv \in E \mid u \in U \wedge w \in W\} .$$



$$U = \{v_1, v_4, v_7\} \quad W = \{v_2, v_3, v_8\} \quad UW = \{v_1v_2, v_7v_8\}$$


Definition 4

For any directed graph (V, E) and any $n \in \mathbb{N}$, a map $p \in E^{[n]}$ is called a **path** in (V, E) of length n iff

$$\forall j, k \in [n] \quad j = k \vee p_j \neq p_k$$
$$\forall j \in [n-1] \exists v, w, x \in V \quad p_j = vw \wedge p_{j+1} = wx .$$

A path is called **simple** iff the nodes along the path are pairwise distinct, except for, possibly, the first and the last node.

Below, **path** shall always mean **simple path**.



Definition 5


For any directed graph (V, E) and any $f \in \mathbb{N}_0^E$, the maps $\varphi^+, \varphi^-, \varphi : 2^V \rightarrow \mathbb{Z}$ such that

$$\forall U \in 2^V \quad \varphi_U^+ = \sum_{uv \in UU^c} f_{uv} \quad (2)$$

$$\varphi_U^- = \sum_{vu \in U^cU} f_{vu} \quad (3)$$

$$\varphi_U = \varphi_U^+ - \varphi_U^- \quad (4)$$

are called the **outflux**, **influx** and **flux** in (V, E) w.r.t. f .



$$U = \{v_1, v_4, v_8\}$$

$$\varphi_U^+ = 3 + 2$$

$$\varphi_U^- = 7 + 5$$

$$\varphi_U = -7$$


For any $u \in V$,

$$\varphi_u^+ := \varphi_{\{u\}}^+$$

$$\varphi_u^- := \varphi_{\{u\}}^-$$

$$\varphi_u := \varphi_{\{u\}}$$

are called the **outflux**, **influx** and **flux** of u in (V, E) w.r.t. f .



$$\varphi_{v_4}^+ = 9$$


$$\varphi_{v_4}^- = 5$$

$$\varphi_{v_4} = 4$$

Lemma 3

For any directed graph (V, E) , any $f \in \mathbb{N}_0^E$ and any $U \subseteq V$

$$\varphi_U = \sum_{u \in U} \varphi_u . \quad (5)$$



Proof.

$$\begin{aligned}\varphi_U &= \sum_{uv \in UU^c} f_{uv} - \sum_{vu \in U^cU} f_{vu} \\ &= \left(\sum_{uv \in UV} f_{uv} - \sum_{uu' \in UU} f_{uu'} \right) - \left(\sum_{vu \in VU} f_{vu} - \sum_{u'u \in UU} f_{u'u'} \right) \\ &= \sum_{uv \in UV} f_{uv} - \sum_{vu \in VU} f_{vu} \\ &= \sum_{u \in U} \left(\sum_{vw \in \{u\}\{u\}^c} f_{vw} - \sum_{vw \in \{u\}^c\{u\}} f_{vw} \right) \\ &= \sum_{u \in U} \varphi_u .\end{aligned}$$

□

Definition 6

A 5-tuple $N = (V, E, s, t, c)$ is called a **network** iff (V, E) is a directed graph and $s \in V$ and $t \in V$ and $s \neq t$ and $c \in \mathbb{N}^E$.

*The nodes s and t are called the **source** and the **sink** of N , respectively.*

*For any edge $e \in E$, c_e is called the **capacity** of e in N .*

Definition 6

A 5-tuple $N = (V, E, s, t, c)$ is called a **network** iff (V, E) is a directed graph and $s \in V$ and $t \in V$ and $s \neq t$ and $c \in \mathbb{N}^E$.

The nodes s and t are called the **source** and the **sink** of N , respectively.

For any edge $e \in E$, c_e is called the **capacity** of e in N .

Definition 6

A 5-tuple $N = (V, E, s, t, c)$ is called a **network** iff (V, E) is a directed graph and $s \in V$ and $t \in V$ and $s \neq t$ and $c \in \mathbb{N}^E$.

The nodes s and t are called the **source** and the **sink** of N , respectively.

For any edge $e \in E$, c_e is called the **capacity** of e in N .

Definition 7

A map $f \in \mathbb{N}_0^E$ is called an **st-preflow** in a network $N = (V, E, s, t, c)$ iff

$$\forall e \in E \quad 0 \leq f_e \leq c_e \quad (6)$$

$$\forall v \in V - \{s\} \quad \varphi_v \leq 0 \ . \quad (7)$$

An *st-preflow* f in N is called an *st-flow* in N iff, in addition,

$$\forall v \in V - \{s, t\} \quad \varphi_v = 0 \ . \quad (8)$$

Definition 7

A map $f \in \mathbb{N}_0^E$ is called an **st-preflow** in a network $N = (V, E, s, t, c)$ iff

$$\forall e \in E \quad 0 \leq f_e \leq c_e \quad (6)$$

$$\forall v \in V - \{s\} \quad \varphi_v \leq 0 . \quad (7)$$

An **st-preflow** f in N is called an **st-flow** in N iff, in addition,

$$\forall v \in V - \{s, t\} \quad \varphi_v = 0 . \quad (8)$$

Definition 8

The instance of the **Maximum *st*-Flow Problem** w.r.t. a network $N = (V, E, s, t, c)$ is to

$$\max_{f \in \mathbb{N}_0^E} \sum_{sv \in E} f_{sv} - \sum_{vs \in E} f_{vs} \quad (9)$$

$$\text{subject to } \forall e \in E \quad 0 \leq f_e \leq c_e \quad (10)$$

$$\forall v \in V - \{s, t\} \quad \sum_{vw \in E} f_{vw} = \sum_{uv \in E} f_{uv} . \quad (11)$$

Note:

$$\sum_{sv \in E} f_{sv} - \sum_{vs \in E} f_{vs} = \varphi_s$$

Definition 9

For any network $N = (V, E, s, t, c)$ and any st -preflow f in N , the **residual network** of N w.r.t. f is the network $N' = (V, E', s, t, c')$ such that

$$E' = E^+ \cup E^-$$

$$E^+ = \{vw \in E \mid c_{vw} - f_{vw} > 0\}$$

$$E^- = \{vw \in V^2 \mid wv \in E \wedge f_{wv} > 0\}$$

and

$$\forall vw \in E' \quad c'_{vw} = \begin{cases} c_{vw} - f_{vw} & \text{if } vw \in E^+ \\ f_{wv} & \text{if } vw \in E^- \end{cases} . \quad (12)$$

For any $e \in E'$, c'_e is called the **residual capacity** of e w.r.t. f .

Any path in (V, E') from s to t (if such a path exists) is called an **augmenting path** of f .

Definition 9

For any network $N = (V, E, s, t, c)$ and any st -preflow f in N , the **residual network** of N w.r.t. f is the network $N' = (V, E', s, t, c')$ such that

$$E' = E^+ \cup E^-$$

$$E^+ = \{vw \in E \mid c_{vw} - f_{vw} > 0\}$$

$$E^- = \{vw \in V^2 \mid wv \in E \wedge f_{wv} > 0\}$$

and

$$\forall vw \in E' \quad c'_{vw} = \begin{cases} c_{vw} - f_{vw} & \text{if } vw \in E^+ \\ f_{wv} & \text{if } vw \in E^- \end{cases} . \quad (12)$$

For any $e \in E'$, c'_e is called the **residual capacity** of e w.r.t. f .

Any path in (V, E') from s to t (if such a path exists) is called an **augmenting path** of f .

Definition 9

For any network $N = (V, E, s, t, c)$ and any st -preflow f in N , the **residual network** of N w.r.t. f is the network $N' = (V, E', s, t, c')$ such that

$$E' = E^+ \cup E^-$$

$$E^+ = \{vw \in E \mid c_{vw} - f_{vw} > 0\}$$

$$E^- = \{vw \in V^2 \mid wv \in E \wedge f_{wv} > 0\}$$

and

$$\forall vw \in E' \quad c'_{vw} = \begin{cases} c_{vw} - f_{vw} & \text{if } vw \in E^+ \\ f_{wv} & \text{if } vw \in E^- \end{cases} . \quad (12)$$

For any $e \in E'$, c'_e is called the **residual capacity** of e w.r.t. f .

Any path in (V, E') from s to t (if such a path exists) is called an **augmenting path** of f .

Lemma 4

Let $N = (V, E, s, t, c)$ be a network and f an st -preflow in N . Assume that an $n \in \mathbb{N}$ and an augmenting path $p = (v_1 w_1, \dots, v_n w_n)$ of f exist.

Let

$$\delta := \min_{vw \in p([n])} c'_{vw} . \quad (13)$$

Then, $f' \in \mathbb{N}_0^E$ such that

$$\forall vw \in E' : \quad f'_{vw} = \begin{cases} f_{vw} + \delta & \text{if } vw \in p([n]) \wedge vw \in E \\ f_{vw} - \delta & \text{if } vw \in p([n]) \wedge wv \in E \\ f_{vw} & \text{otherwise} \end{cases} \quad (14)$$

is an st -preflow in N w.r.t. which

$$\varphi'_s = \varphi_s + \delta . \quad (15)$$

Moreover, if f is an st -flow in N , so is f' .

Lemma 4

Let $N = (V, E, s, t, c)$ be a network and f an st -preflow in N . Assume that an $n \in \mathbb{N}$ and an augmenting path $p = (v_1 w_1, \dots, v_n w_n)$ of f exist. Let

$$\delta := \min_{vw \in p([n])} c'_{vw} . \quad (13)$$

Then, $f' \in \mathbb{N}_0^E$ such that

$$\forall vw \in E' : \quad f'_{vw} = \begin{cases} f_{vw} + \delta & \text{if } vw \in p([n]) \wedge vw \in E \\ f_{vw} - \delta & \text{if } vw \in p([n]) \wedge wv \in E \\ f_{vw} & \text{otherwise} \end{cases} \quad (14)$$

is an st -preflow in N w.r.t. which

$$\varphi'_s = \varphi_s + \delta . \quad (15)$$

Moreover, if f is an st -flow in N , so is f' .

Lemma 4

Let $N = (V, E, s, t, c)$ be a network and f an st -preflow in N . Assume that an $n \in \mathbb{N}$ and an augmenting path $p = (v_1 w_1, \dots, v_n w_n)$ of f exist. Let

$$\delta := \min_{vw \in p([n])} c'_{vw} . \quad (13)$$

Then, $f' \in \mathbb{N}_0^E$ such that

$$\forall vw \in E' : \quad f'_{vw} = \begin{cases} f_{vw} + \delta & \text{if } vw \in p([n]) \wedge vw \in E \\ f_{vw} - \delta & \text{if } vw \in p([n]) \wedge wv \in E \\ f_{vw} & \text{otherwise} \end{cases} \quad (14)$$

is an st -preflow in N w.r.t. which

$$\varphi'_s = \varphi_s + \delta . \quad (15)$$

Moreover, if f is an st -flow in N , so is f' .

Lemma 4

Let $N = (V, E, s, t, c)$ be a network and f an st -preflow in N . Assume that an $n \in \mathbb{N}$ and an augmenting path $p = (v_1 w_1, \dots, v_n w_n)$ of f exist. Let

$$\delta := \min_{vw \in p([n])} c'_{vw} . \quad (13)$$

Then, $f' \in \mathbb{N}_0^E$ such that

$$\forall vw \in E' : \quad f'_{vw} = \begin{cases} f_{vw} + \delta & \text{if } vw \in p([n]) \wedge vw \in E \\ f_{vw} - \delta & \text{if } vw \in p([n]) \wedge wv \in E \\ f_{vw} & \text{otherwise} \end{cases} \quad (14)$$

is an st -preflow in N w.r.t. which


$$\varphi'_s = \varphi_s + \delta . \quad (15)$$

Moreover, if f is an st -flow in N , so is f' .

Definition 10

Let (V, E) be a directed graph. Let $s \in V$ and $t \in V$ and $s \neq t$.

- ▶ $X \subseteq V$ is called an ***st-cutset*** of (V, E) iff $s \in X$ and $t \notin X$.
- ▶ $Y \subseteq E$ is called an ***st-cut*** of (V, E) iff there exists an *st-cutset* X such that $Y = \{vw \in E \mid v \in X \wedge w \notin X\}$.



Definition 11

The instance of the **Minimum *st*-Cut Problem** w.r.t. a network $N = (V, E, s, t, c)$ is to

$$\min_{x \in \{0,1\}^V} \sum_{vw \in E} x_v(1 - x_w)c_{vw} \quad (16)$$

$$\text{subject to } x_s = 1 \quad (17)$$

$$x_t = 0 \quad (18)$$

Note: With $X := \{v \in V | x_v = 1\}$, we have

$$\sum_{vw \in E} x_v(1 - x_w)c_{vw} = \sum_{vw \in XX^c} c_{vw}$$

Lemma 5

For every network $N = (V, E, s, t, c)$, every st -flow f in N , and every st -cutset $X \subseteq V$,

$$\varphi_s \leq \sum_{vw \in XX^c} c_{vw} . \quad (19)$$

Proof.

$$\begin{aligned}\varphi_s &= \sum_{v \in S} \varphi_v && \text{by (8) and } t \notin S \\ &= \varphi_S && \text{by Lemma 3} \\ &\leq \varphi_S^+ && \text{by (3), (4) and } 0 \leq f \\ &= \sum_{vw \in SS^c} f_{vw} && \text{by (2)} \\ &\leq \sum_{vw \in SS^c} c_{vw} && \text{by (6).}\end{aligned}$$

□

Lemma 5 does **not** hold analogously for every st -preflow, because, w.r.t. an st -preflow, φ_S need not be an upper bound on φ_s .

Theorem 1

For any network $N = (V, E, s, t, c)$, any $s, t \in V$ such that $s \neq t$, and any st -flow f in N , the following three conditions are equivalent

- 1. There exists an st -cut whose capacity is equal to φ_s .*
- 2. The st -flow f is optimal, i.e., a solution of (9)–(11).*
- 3. No augmenting path of f exists.*

Proof.

(1) implies (2) by virtue of Lemma 5.

(2) implies (3) by virtue of Lemma 4.

We prove that (3) implies (1):

- ▶ Let f be an st -flow such that no augmenting path exists.
- ▶ Let S be the set of all nodes $v \in V$ such that there exists a path in the residual network w.r.t. f from s to v . Let S also include s itself.
- ▶ Then, $t \notin S$ (otherwise, the path from s to t in the residual network would be an augmenting path).
- ▶ Moreover, ...

► Moreover,

$$\begin{aligned}\varphi_s &= \sum_{v \in S} \varphi_v && \text{by (8) and } t \notin S \\ &= \varphi_S && \text{by Lemma 3} \\ &= \sum_{vw \in SS^c} f_{vw} - \sum_{vw \in S^c S} f_{vw} && \text{by definition of } \varphi_S \\ &= \sum_{vw \in SS^c} c_{vw} && \text{by the arguments below.}\end{aligned}$$

- For any $vw \in SS^c$, we have $f_{vw} = c_{vw}$ (otherwise, the contradiction $w \in S$ follows by construction of S and by definition of the residual network).
- For any $vw \in S^c S$, we have $f_{vw} = 0$ (otherwise, the contradiction $v \in S$ follows by construction of S and by definition of the residual network).

□

Algorithm 1 (Ford and Fulkerson, 1956)

Input: Network $N = (V, E, s, t, c)$

Output: $f : E \rightarrow \mathbb{N}_0$

for all $vw \in E$

$$f_{vw} := 0$$

while $\exists n \in \mathbb{N} \exists$ augmenting path $p = (v_1 w_1, \dots, v_n w_n)$ of f

$$\delta := \min_{vw \in p([n])} c'_{vw}$$

for all $vw \in E$

$$f_{vw} := \begin{cases} f_{vw} + \delta & \text{if } vw \in P \wedge vw \in E \\ f_{vw} - \delta & \text{if } vw \in P \wedge wv \in E \\ f_{vw} & \text{otherwise} \end{cases}$$

Theorem 2

Algorithm 1 terminates. The output f is a maximum st -flow in N .

Proof. Termination.

- ▶ For every augmenting path processed, φ_s increases by at least 1.
- ▶ Moreover,

$$\varphi_s \leq \sum_{vw \in \{s\}\{s\}^c} c_{vw} \quad (\text{by Lemma 5})$$

- ▶ Therefore, only finitely many augmenting paths are processed.
- ▶ Thus, the algorithm terminates.

Optimality:


- ▶ Throughout the execution, f is an st -flow in N .
- ▶ When the algorithm terminates, no augmenting path exists.
- ▶ Thus, f is a maximum st -flow in N (by Theorem 1).

Note: An implementation with runtime complexity $O(|E|\varphi_s)$ exists.

Definition 12

For any $n \in \mathbb{N}$ and any $c \in C_{n2}^+$, the **network** $N = (V, E, s, t, w)$ of c contains the nodes $V = \{s, t, 1, \bar{1}, \dots, n, \bar{n}\}$ and the weighted edges

for any $c_{\{j\}\emptyset} > 0$	$s\bar{j}, j\bar{t}$	$w_{s\bar{j}} := w_{j\bar{t}} := \frac{1}{2}c_{\{j\}\emptyset}$
for any $c_{\emptyset\{j\}} > 0$	$s\bar{j}, \bar{j}t$	$w_{s\bar{j}} := w_{\bar{j}t} := \frac{1}{2}c_{\emptyset\{j\}}$
for any $c_{\{j,k\}\emptyset} > 0$	$j\bar{k}, k\bar{j}$	$w_{j\bar{k}} := w_{k\bar{j}} := \frac{1}{2}c_{\{j,k\}\emptyset}$
for any $c_{\{j\}\{k\}} > 0$	$j\bar{k}, \bar{k}\bar{j}$	$w_{j\bar{k}} := w_{\bar{k}\bar{j}} := \frac{1}{2}c_{\{j\}\{k\}}$
for any $c_{\emptyset\{j,k\}} > 0$	$\bar{j}k, \bar{k}j$	$w_{\bar{j}k} := w_{\bar{k}j} := \frac{1}{2}c_{\emptyset\{j,k\}}$



Definition 13

For any $n \in \mathbb{N}$, any $c \in C_{n2}^+$, the network $N = (V, E, s, t, w)$ of c and any $x \in \{0, 1\}^n$, let $x' \in \{0, 1\}^V$ such that

$$x'_s = 1 \tag{20}$$

$$x'_t = 0 \tag{21}$$

$$\forall j \in [n] \quad x'_j = x_j \tag{22}$$

$$\forall j \in [n] \quad x'_{\bar{j}} = 1 - x_j \tag{23}$$

Lemma 6

For any $n \in \mathbb{N}$, any $c \in C_{n2}^+$, the network $N = (V, E, s, t, w)$ of c and any $x \in \{0, 1\}^n$, the PBF $f_c : \{0, 1\}^n \rightarrow \mathbb{R}$ defined by c is such that

$$\forall x \in \{0, 1\}^n \quad f_c(x) = \sum_{jk \in E} w_{jk} x'_j (1 - x'_k) . \quad (24)$$


Proof. Trivial.

Lemma 7


For any $n \in \mathbb{N}$, any $c \in C_{n2}^+$, the network $N = (V, E, s, t, w)$ of c , any $m \in \mathbb{N}$ and any augmenting path (sv_1, \dots, v_mt) in N , the path $(s\bar{v}_m, \dots, \bar{v}_1t)$ is also augmenting.

Example.

Given path $sj, j\bar{k}, \bar{k}\bar{j}, \bar{j}t$



Conjectured path $sj, jk, k\bar{j}, \bar{j}t$



Posiform corresponding to both paths

$$(1 - x_j) + x_j x_k + (1 - x_k) x_j + (1 - x_j)$$

Lemma 8

For any $n \in \mathbb{N}$, any $c \in C_{n2}^+$, the network $N = (V, E, s, t, w)$ of c , any st -flow $g \in \mathbb{R}^E$ in N , and the posiform $c' \in C_{n2}^+$ of the residual network of g ,

$$f_c = c_{\emptyset\emptyset} + \varphi_s + f_{c'} . \quad (25)$$

Moreover, the r.h.s. is a posiform which differs from the homogenous posiform c' only by the additional constant term $c_{\emptyset\emptyset} + \varphi_s$.

Lemma 9

For any $n \in \mathbb{N}$, any $f : \{0, 1\}^n \rightarrow \mathbb{R}$ and any $c, c' \in C_{n2}^+(f)$ such that $c_{\emptyset\emptyset} < c'_{\emptyset\emptyset}$, there exists an augmenting path in the network of c .

Theorem 3

For any $n \in \mathbb{N}$, any $f : \{0, 1\}^n \rightarrow \mathbb{R}$, any $c \in C_{n2}^+(f)$ and the value $\hat{\varphi}_s \in \mathbb{R}$ of any maximum st -flow in the network of c ,

$$r_f = c_{\emptyset\emptyset} + \hat{\varphi}_s . \quad (26)$$

Summary:

- ▶ The computation of the floor dual has been reduced to the Maximum *st*-Flow Problem.


Theorem 4 (strong persistency)

For any $n \in \mathbb{N}$, any $f : \{0, 1\}^n \rightarrow \mathbb{R}$, any $c \in C_{n2}^+(f)$, any maximum st -flow $g \in \mathbb{R}^E$ in the network of c and the set $S \subseteq V$ of all nodes reachable from s via a path in the residual network of g ,

$$\forall \hat{x} \in \operatorname{argmin}_{x \in \{0,1\}^n} f(x) \quad \forall j \in [n] \quad (j \in S \Rightarrow x_j = 1) \wedge (\bar{j} \in S \Rightarrow x_j = 0) .$$

Proof.


- ▶ Let $c' \in C_{n_2}^+$ be the posiform of the residual network of g .
- ▶ Firstly, $c'_{\emptyset\emptyset} = 0$ (by Definition 12)
- ▶ Secondly, $\forall j \in [n] : j \notin S \vee \bar{j} \notin S$ by the following argument. If $j \in S \wedge \bar{j} \in S$, there exist paths in the residual network



in contradiction to the maximality of the flow g .

Proof.


- ▶ Let $c' \in C_{n2}^+$ be the posiform of the residual network of g .
- ▶ Firstly, $c'_{\emptyset\emptyset} = 0$ (by Definition 12)
- ▶ Secondly, $\forall j \in [n] : j \notin S \vee \bar{j} \notin S$ by the following argument. If $j \in S \wedge \bar{j} \in S$, there exist paths in the residual network



in contradiction to the maximality of the flow g .

Proof.

- ▶ Let $c' \in C_{n2}^+$ be the posiform of the residual network of g .
- ▶ Firstly, $c'_{\emptyset\emptyset} = 0$ (by Definition 12)
- ▶ Secondly, $\forall j \in [n] : j \notin S \vee \bar{j} \notin S$ by the following argument. If $j \in S \wedge \bar{j} \in S$, there exist paths in the residual network



in contradiction to the maximality of the flow g .

► Thirdly,

$$c'_{\{j,k\}\emptyset} > 0 \wedge j \in S \Rightarrow \bar{k} \in S$$


$$c'_{\{j\}\{k\}} > 0 \wedge j \in S \Rightarrow k \in S$$

$$c'_{\{j\}\{k\}} > 0 \wedge \bar{k} \in S \Rightarrow \bar{j} \in S$$

$$c'_{\emptyset\{j,k\}} > 0 \wedge \bar{j} \in S \Rightarrow k \in S$$

(because edges with positive weight cannot direct from a node in S to a node not in S , by definition of S).

Let $c'_{\{j,k\}\emptyset} > 0 \wedge j \in S$.




► Thirdly,

$$\begin{aligned}c'_{\{j,k\}\emptyset} > 0 \wedge j \in S &\Rightarrow \bar{k} \in S \\c'_{\{j\}\{k\}} > 0 \wedge j \in S &\Rightarrow k \in S \\c'_{\{j\}\{k\}} > 0 \wedge \bar{k} \in S &\Rightarrow \bar{j} \in S \\c'_{\emptyset\{j,k\}} > 0 \wedge \bar{j} \in S &\Rightarrow k \in S\end{aligned}$$

(because edges with positive weight cannot direct from a node in S to a node not in S , by definition of S).

Let $c'_{\{j\}\{k\}} > 0 \wedge j \in S$.




► Thirdly,

$$\begin{aligned}c'_{\{j,k\}\emptyset} > 0 \wedge j \in S &\Rightarrow \bar{k} \in S \\c'_{\{j\}\{k\}} > 0 \wedge j \in S &\Rightarrow k \in S \\c'_{\{j\}\{k\}} > 0 \wedge \bar{k} \in S &\Rightarrow \bar{j} \in S \\c'_{\emptyset\{j,k\}} > 0 \wedge \bar{j} \in S &\Rightarrow k \in S\end{aligned}$$

(because edges with positive weight cannot direct from a node in S to a node not in S , by definition of S).

$$\text{Let } c'_{\{j\}\{k\}} > 0 \wedge \bar{k} \in S.$$



► Thirdly,

$$c'_{\{j,k\}\emptyset} > 0 \wedge j \in S \Rightarrow \bar{k} \in S$$


$$c'_{\{j\}\{k\}} > 0 \wedge j \in S \Rightarrow k \in S$$

$$c'_{\{j\}\{k\}} > 0 \wedge \bar{k} \in S \Rightarrow \bar{j} \in S$$

$$c'_{\emptyset\{j,k\}} > 0 \wedge \bar{j} \in S \Rightarrow k \in S$$

(because edges with positive weight cannot direct from a node in S to a node not in S , by definition of S).

Let $c'_{\emptyset\{j,k\}} > 0 \wedge \bar{j} \in S$.



- Therefore, (S', y) with

$$S' = \{j \in [n] \mid j \in S \vee \bar{j} \in S\}$$

and $y : S' \rightarrow \{0, 1\}$ such that

$$\forall j \in S' \quad y_j = \begin{cases} 1 & \text{if } j \in S \\ 0 & \text{if } \bar{j} \in S \end{cases}$$

is a **contractor** of c' .

- Fourthly, **weak persistency** holds for (S, y) at the minima of f (because $r_f + f_{c'} = f$).

- Therefore, (S', y) with

$$S' = \{j \in [n] \mid j \in S \vee \bar{j} \in S\}$$

and $y : S' \rightarrow \{0, 1\}$ such that

$$\forall j \in S' \quad y_j = \begin{cases} 1 & \text{if } j \in S \\ 0 & \text{if } \bar{j} \in S \end{cases}$$

is a **contractor** of c' .

- Fourthly, **weak persistency** holds for (S, y) at the minima of f (because $r_f + f_{c'} = f$).

► Moreover, $\forall x \in \{0, 1\}^n$:

$$\begin{aligned} f_{c'}(x) &= \sum_{jk \in SS} w_{jk} x'_j (1 - x'_k) + \sum_{jk \in SSC} w_{jk} x'_j (1 - x'_k) \\ &+ \sum_{jk \in S^C S} w_{jk} x'_j (1 - x'_k) + \sum_{jk \in S^C S^C} w_{jk} x'_j (1 - x'_k) \end{aligned}$$

If $\forall j \in S' : x_j = y_j$,

$$f_{c'}(x) = \sum_{jk \in S^C S^C} w_{jk} x'_j (1 - x'_k)$$

Otherwise,

$$f_{c'}(x) > \sum_{jk \in S^C S^C} w_{jk} x'_j (1 - x'_k)$$

Thus, strong persistency holds.